

Supplementary material has been published as submitted. It has not been copyedited, or typeset by Acta Oncologica

Table S1. Contouring of the CTVe according to risk-adapted NOAC guidelines

	Clinical staging ¹	Deauville score 4-5 ²	Deauville score 3-5 ²
	<i>n</i> (%)	<i>n</i> (%)	<i>n</i> (%)
Cranial border of CTV			
Very low	32 (19.3)	32 (20.1)	27 (17.0)
Low	14 (8.4)	15 (9.4)	15 (9.4)
Intermediate	99 (59.6)	93 (58.5)	93 (58.5)
High	21 (12.7)	19 (11.9)	24 (15.1)
Inclusion of external iliac region			
Yes	98 (59.0)	93 (58.5)	98 (61.6)
No	68 (41.0)	66 (41.5)	61 (38.4)

¹All patients (*n* = 166)

²Assessed for 159 patients with PET-CT available

Table S2. Age differences between clinical staging and retrospective PET-CT assessment

	Clinical staging ¹			Deauville score 4-5 ²			Deauville Score 3-5 ²		
	Age, <i>n</i> (%)		<i>P</i> ³	Age, <i>n</i> (%)		<i>P</i> ³	Age, <i>n</i> (%)		<i>P</i> ³
	< 70 years	≥ 70 years		< 70 years	≥ 70 years		< 70 years	≥ 70 years	
	120 (72.3)	46 (27.7)		115 (72.3)	44 (27.7)		115 (72.3)	44 (27.7)	
Cranial border			0.14			0.65			0.53
Very low or Low	29 (24.2)	17 (37.0)		32 (27.8)	15 (34.1)		28 (24.3)	14 (31.8)	
Intermediate	73 (60.8)	26 (56.5)		68 (59.1)	25 (56.8)		68 (59.1)	25 (56.8)	
High	18 (15.0)	3 (6.5)		15 (13.0)	4 (9.1)		19 (16.5)	5 (11.4)	
Inclusion of the external iliac region			0.001			0.015			0.009
Yes	80 (66.7)	18 (39.1)		74 (64.3)	19 (43.2)		78 (67.8)	20 (45.5)	
No	40 (33.3)	28 (60.9)		41 (35.7)	25 (56.8)		37 (32.2)	24 (54.5)	

¹ All patients (*n* = 166)² Assessed for 159 patients with PET-CT available³ Using the chi-squared test