

- S1. Abraham A, Hart A, Andrade I, Hackney ME. Dynamic Neuro-Cognitive Imagery Improves Mental Imagery Ability, Disease Severity, and Motor and Cognitive Functions in People with Parkinson's Disease. *Neural Plast* 2018; : 6168507. DOI: 10.1155/2018/6168507
- S2. Acarer A, Karapolat H, Celebisoy N, Ozgen G, Colakoglu Z. Is customized vestibular rehabilitation effective in patients with Parkinson's? *NeuroRehabilitation* 2015; 37(2): 255–262. DOI: 10.3233/NRE-151258
- S3. Albrecht F, Pereira JB, Mijalkov M, Freidle M, Johansson H, Ekman U, et al. Effects of a Highly Challenging Balance Training Program on Motor Function and Brain Structure in Parkinson's Disease. *J Parkinsons Dis* 2021; 11(4): 2057–2071. DOI: 10.3233/JPD-212801
- S4. Alencar de Lima T, Ferreira-Moraes R, Gomes da Costa Alves WM, Gonçalves Gibson Alves T, Pantoja Pimentel C. Resistance training reduces depressive symptoms in elderly people with Parkinson disease: a controlled randomized study. *Scand J Med Sci Sports* 2019; 29(12): 1957–1967. DOI: 10.1111/sms.13528
- S5. Arfa-Fatollahkhani P, Safar Cherati A, Habibi SAH, Shahidi GA, Sohrabi A, Zamani B. Effects of treadmill training on the balance, functional capacity and quality of life in Parkinson's disease: A randomized clinical trial. *J Complement Integr Med [Serial on the Internet]* 2020; 17(1): 1–9. DOI: 10.1515/jcim-2018-0245
- S6. Ashburn A, Fazakarley L, Ballinger C, Pickering R, McLellan LD, Fitton C, et al. A randomised controlled trial of a home based exercise programme to reduce the risk of falling among people with Parkinson's disease. *J Neurol Neurosurgery, Psychiatry* 2007 Jul;78(7):678-684 2007; 78(7): 678–684. DOI: 10.1136/jnnp.2006.099333
- S7. Atan T, Ozyemisci Taskiran O, Bora Tokcaer A, Kaymak Karatas G, Karakus Caliskan A, Karaoglan B. Effects of different percentages of body weight-supported treadmill training in Parkinson's disease: a double-blind randomized controlled trial. *Turkish J Med Sci* 2019; 49(4): 999–1007. DOI: 10.3906/sag-1812-57
- S8. Cakit BD, Saracoglu MM, Genc H, Erdem HR, Inan L, L CBSMGHEHI. The effects of incremental speed-dependent treadmill training on postural instability and fear of failing in Parkinson's disease. *Clin Rehabil* 2007; 21(8): 698–705. DOI: 10.1177/0269215507077269
- S9. Capato TTC, de Vries NM, Int'Hout J, Barbosa ER, Nonnekes J, Bloem BR. Multimodal Balance Training Supported by Rhythmical Auditory Stimuli in Parkinson's Disease: A Randomized Clinical Trial. *J Park Dis* 2020; 10(1): 333–346. DOI: 10.3233/JPD-191752
- S10. Capecci M, Pournajaf S, Galafate D, Sale P, Le Pera D, Goffredo M, et al. Clinical effects of robot-assisted gait training and treadmill training for Parkinson's disease. A randomized controlled trial. *Ann Phys Rehabil Med* 2019 Sep;62(5):303-312 2019; 62(5): 303–312. DOI: 10.1016/j.rehab.2019.06.016
- S11. Carda S, Invernizzi M, Baricich A, Comi C, Croquelois A, Cisari C. Robotic gait training is not superior to conventional treadmill training in parkinson disease: a single-blind randomized controlled trial. *Neurorehabil Neural Repair* 2012; 26(9): 1027–1034. DOI: 10.1177/1545968312446753
- S12. Cheng F-Y, Yang Y-R, Wu Y-R, Cheng S-J, Wang R-Y. Effects of curved-walking training on curved-walking performance and freezing of gait in individuals with Parkinson's disease: A randomized controlled trial. *Parkinsonism Relat Disord* 2017; 43: 20–26. DOI: 10.1016/j.parkreldis.2017.06.021
- S13. Cherup NP, Strand KL, Lucchi L, Wooten S V, Luca C, Signorile JF. Yoga Meditation Enhances Proprioception and Balance in Individuals Diagnosed With Parkinson's Disease. *Percept Mot Skills* 2021; 128(1): 304–323. DOI: 10.1177/0031512520945085
- S14. Choi H-J, Garber CE, Jun T-W, Jin Y-S, Chung S-J, Kang H-J. Therapeutic effects of tai chi in patients with Parkinson's disease. *ISRN Neurol* 2013; 2013: 548240. DOI: 10.1155/2013/548240
- S15. Collett J, Franssen M, Meaney A, Wade D, Izadi H, Tims M, et al. Phase II randomised controlled trial of a 6-month self-managed community exercise programme for people with Parkinson's disease. *J Neurol Neurosurg psychiatry* 2016; 88(3): 204–211. DOI: 10.1136/jnnp-2016-314508
- S16. Combs SA, Diehl MD, Chrzastowski C, Didrick N, McCoin B, Mox N, et al. Community-based group exercise for persons with Parkinson disease: a randomized controlled trial. *NeuroRehabilitation* 2013; 32(1): 117–124. DOI: 10.3233/NRE-130828
- S17. Conradsson D, Löfgren N, Nero H, Hagströmer M, Ståhle A, Lökk J, et al. The Effects of Highly Challenging Balance Training in Elderly With Parkinson's Disease: A Randomized Controlled Trial. *Neurorehabil Neural Repair* 2015; 29(9): 827–836. DOI: 10.1177/1545968314567150
- S18. Cugusi L, Solla P, Serpe R, Carzedda T, Piras L, Oggianu M, et al. Effects of a Nordic Walking program on motor and non-motor symptoms, functional performance and body composition in patients with Parkinson's disease. *NeuroRehabilitation* 2015; 37(2): 245–254. DOI: 10.3233/NRE-151257
- S19. da Silva AZ, Israel VL. Effects of dual-task aquatic exercises on functional mobility, balance and gait of individuals with Parkinson's disease: A randomized clinical trial with a 3-month follow-up. *Complement Ther Med* 2019; 42: 119–124. DOI: 10.1016/j.ctim.2018.10.023
- S20. Dipasquale S, Meroni R, Sasanelli F, Messineo I, Piscitelli D, Perin C, et al. Physical Therapy Versus a General Exercise Programme in Patients with Hoehn Yahr Stage II Parkinson's Disease: A Randomized Controlled Trial. *J Parkinsons Dis* 2017; 7(1): 203–210. DOI: 10.3233/JPD-161015

- S21. Ebersbach G, Ebersbach A, Edler D, Kaufhold O, Kusch M, Kupsch A, et al. Comparing exercise in Parkinson's disease -- the Berlin LSVTBIG study. *Mov Disord* 2010; 25(12): 1902-1908. DOI: 10.1002/mds.23212
- S22. Ganesan M, Sathyaprabha TN, Gupta A, Pal PK. Effect of partial weight-supported treadmill gait training on balance in patients with Parkinson disease. *PM&R* 2014; 6(1): 22-33. DOI: 10.1016/j.pmrj.2013.08.604
- S23. Gao Q, Leung A, Yang Y, Wei Q, Guan M, Jia C, et al. Effects of Tai Chi on balance and fall prevention in Parkinson's disease: a randomized controlled trial. *Clin Rehabil* 2014; 28(8): 748-753. DOI: 10.1177/0269215514521044
- S24. Goodwin VA, Richards SH, Henley W, Ewings P, Taylor AH, Campbell JL, et al. An exercise intervention to prevent falls in people with Parkinson's disease: a pragmatic randomised controlled trial. *J Neurol Neurosurgery, Psychiatry* 2011 Nov;82(11):1232-1238 2011; 82(11): 1232-1238. DOI: 10.1136/jnnp-2011-300919
- S25. Hackney ME, Earhart GM. EFFECTS OF DANCE ON MOVEMENT CONTROL IN PARKINSON'S DISEASE: A COMPARISON OF ARGENTINE TANGO AND AMERICAN BALLROOM. *J Rehabil Med* 2009; 41(6): 475-481. DOI: 10.2340/16501977-0362
- S26. Hubble RP, Naughton G, Silburn PA, Cole MH. Trunk Exercises Improve Gait Symmetry in Parkinson Disease: A Blind Phase II Randomized Controlled Trial. *Am J Phys Med Rehabil* 2018; 97(3): 151-159. DOI: 10.1097/PHM.0000000000000858
- S27. Johansson H, Freidle M, Ekman U, Schalling E, Leavy B, Svenningsson P, et al. Feasibility Aspects of Exploring Exercise-Induced Neuroplasticity in Parkinson's Disease: A Pilot Randomized Controlled Trial. *Park Dis* 2020 Mar 25;(2410863)Epub 2020; 2020: 2410863. DOI: 10.1155/2020/2410863
- S28. Jung SH, Hasegawa N, Mancini M, King LA, Carlson-Kuhta P, Smulders K, et al. Effects of the agility boot camp with cognitive challenge (ABC-C) exercise program for Parkinson's disease. *NPJ Park Dis* 2020 Nov 2;6(31)Epub 2020; 6(1): 31. DOI: 10.1038/s41531-020-00132-z
- S29. Khalil H, Busse M, Quinn L, Nazzal M, Batyha W, Alkhazaleh S, et al. A pilot study of a minimally supervised home exercise and walking program for people with Parkinson's disease in Jordan. *Neurodegener Dis Manag* 2017; 7(1): 73-84. DOI: 10.2217/nmt-2016-0041
- S30. Khuzema A, Brammatha A, Arul Selvan V. Effect of home-based Tai Chi, Yoga or conventional balance exercise on functional balance and mobility among persons with idiopathic Parkinson's disease: An experimental study. *Hong Kong Physiother J* 2020; 40(1): 39-49. DOI: 10.1142/S1013702520500055
- S31. Kurt EE, Büyükturan B, Büyükturan Ö, Erdem HR, Tuncay F. Effects of Ai Chi on balance, quality of life, functional mobility, and motor impairment in patients with Parkinson's disease<sup/>. *Disabil Rehabil* 2018; 40(7): 791-797. DOI: 10.1080/09638288.2016.1276972
- S32. Kwok JYY, Kwan JCY, Auyeung M, Mok VCT, Lau CKY, Choi KC, et al. Effects of mindfulness yoga versus stretching and resistance training exercises on anxiety and depression for people with Parkinson disease: a randomized clinical trial [with consumer summary]. *JAMA Neurol* 2019 Jul;76(7):755-763 2019; 76(7): 755-763. DOI: 10.1001/jamaneurol.2019.0534
- S33. Landers MR, Hatlevig RM, Davis AD, Richards AR, Rosenlof LE. Does attentional focus during balance training in people with Parkinson's disease affect outcome? A randomised controlled clinical trial. *Clin Rehabil* 2016; 30(1): 53-63. DOI: 10.1177/0269215515570377
- S34. Leal LC, Abrahim O, Rodrigues RP, da Silva MC, Araújo AP, de Sousa EC, et al. Low-volume resistance training improves the functional capacity of older individuals with Parkinson's disease. *Geriatr Gerontol Int* 2019; 19(7): 635-640. DOI: 10.1111/ggi.13682
- S35. Lee N-Y, Lee D-K, Song H-S, H-S LN-YLD-KS. Effect of virtual reality dance exercise on the balance, activities of daily living, and depressive disorder status of Parkinson's disease patients. *J Phys Ther Sci* 2015 Jan;27(1):145-147 2015; 27(1): 145-147. DOI: 10.1589/jpts.27.145
- S36. Lee HJ, Kim SY, Chae Y, Kim MY, Yin C, Jung WS, et al. Turo (Qi Dance) Program for Parkinson's Disease Patients: randomized, Assessor Blind, Waiting-List Control, Partial Crossover Study. *Explore (NY) [Serial on the Internet]* 2018; 14(3): 216-223. DOI: 10.1016/j.explore.2017.11.002
- S37. Liu XL, Chen S, Wang Y. Effects of Health Qigong Exercises on Relieving Symptoms of Parkinson's Disease. *Evid Based Complement Alternat Med* 2016; 2016: 5935782. DOI: 10.1155/2016/5935782
- S38. Li F, Harmer P, Fitzgerald K, Eckstrom E, Stock R, Galver J, et al. Tai chi and postural stability in patients with Parkinson's disease. *N Engl J Med* 2012; 366(6): 511-519. DOI: 10.1056/NEJMoa1107911
- S39. Li G, Huang P, Cui SS, Tan YY, He YC, Shen X, et al. Mechanisms of motor symptom improvement by long-term Tai Chi training in Parkinson's disease patients. *Transl Neurodegener [Serial on the Internet]* 2022 [cited 2022 May 26]; 11(1): 1-10. DOI: 10.1186/S40035-022-00280-7/FIGURES/2
- S40. Liao Y-Y, Yang Y-R, Cheng S-J, Wu Y-R, Fuh J-L, Wang R-Y. Virtual Reality-Based Training to Improve Obstacle-Crossing Performance and Dynamic Balance in Patients With Parkinson's Disease. *Neurorehabil Neural Repair* 2015; 29(7): 658-667. DOI: 10.1177/1545968314562111
- S41. Mak MKY, Wong-Yu ISK. Six-Month Community-Based Brisk Walking and Balance Exercise Alleviates Motor Symptoms and Promotes Functions in People with Parkinson's Disease: a Randomized Controlled Trial. *J Parkinsons Dis [Serial on the Internet]* 2021; . DOI: 10.3233/JPD-202503
- S42. Mollinedo-Cardalda I, Cancela-Carral JM, Vila-Suárez MH. Effect of a Mat Pilates Program with TheraBand on Dynamic Balance in Patients with Parkinson's Disease: Feasibility Study and Randomized Controlled Trial. *Rejuvenation Res* 2018; 21(5): 423-430. DOI: 10.1089/rej.2017.2007
- S43. Monticone M, Ambrosini E, Laurini A, Rocca B, Foti C. In-patient multidisciplinary rehabilitation for Parkinson's disease: A randomized controlled trial. *Mov Disord* 2015; 30(8): 1050-1058. DOI: 10.1002/mds.26256

- S44. Moon J-H, Jung J-H, Cho H-Y. Effects of balance training using a wii fit balance board on balance, gait and activities of daily living in patients with parkinson disease: a pilot, randomized controlled trial. *Medico-legal Updat* [Serial on the Internet] 2020; 20(1): 1799-1803. DOI: 10.37506/v20/il/2020/mlu/194564
- S45. Morris ME, Iansek R, Kirkwood B, B MMIRK, Morris ME, Iansek R, et al. A randomized controlled trial of movement strategies compared with exercise for people with Parkinson's disease. *Mov Disord* 2009; 24(1): 64–71. DOI: 10.1002/mds.22295
- S46. Morris ME, Menz HB, McGinley JL, Watts JJ, Huxham FE, Murphy AT, et al. A Randomized Controlled Trial to Reduce Falls in People With Parkinson's Disease. *Neurorehabil Neural Repair* 2015; 29(8): 777-785. DOI: 10.1177/1545968314565511
- S47. Nadeau A, Pourcher E, Corbeil P. Effects of 24 wk of treadmill training on gait performance in Parkinson's disease. *Med Sci Sports Exerc* 2014; 46(4): 645–655. DOI: 10.1249/MSS.0000000000000144
- S48. Ni M, Signorile JF, Mooney K, Balachandran A, Potiaumpai M, Luca C, et al. Comparative effect of power training and high-speed yoga on motor function in older patients with Parkinson disease. *Arch Phys Med Rehabil* 2016 Mar;97(3)345-354 2016; 97(3): 345-354.e15. DOI: 10.1016/j.apmr.2015.10.095
- S49. Nieuwboer A, Kwakkel G, Rochester L, Jones D, van Wegen E, Willems AM, et al. Cueing training in the home improves gait-related mobility in Parkinson's disease: the RESCUE trial. *J Neurol Neurosurg Psychiatry* 2007; 78(2): 134–140. DOI: 10.1136/jnnp.200X.097923
- S50. Ortiz-Rubio A, Cabrera-Martos I, Torres-Sánchez I, Casilda-López J, López-López L, Valenza MC. Effects of a resistance training program on balance and fatigue perception in patients with Parkinson's disease: A randomized controlled trial. *Med Clin (Barc)* 2018; 150(12): 460–464. DOI: 10.1016/j.medcli.2017.10.022
- S51. Paul SS, Canning CG, Song J, Fung VSC, Sherrington C, C PSCCSJFVS, et al. Leg muscle power is enhanced by training in people with Parkinson's disease: a randomized controlled trial [with consumer summary]. *Clin Rehabil* 2014 Mar;28(3)275-288 2014; 28(3): 275–288. DOI: 10.1177/0269215513507462
- S52. de la Cruz S, S P de la C, Pérez de la Cruz S. Effectiveness of aquatic therapy for the control of pain and increased functionality in people with Parkinson's disease: a randomized clinical trial. *Eur J Phys Rehabil Med* 2017; 53(6): 825–832. DOI: 10.23736/S1973-9087.17.04647-0
- S53. Perez-de la Cruz S, S P la C, Pérez-de la Cruz S, S P la C, la Cruz S. A bicentric controlled study on the effects of aquatic Ai Chi in Parkinson disease. *Complement Ther Med* [Serial on the Internet] 2018; 36(pp 147-153 CC-Complementary Medicine): 147-153. DOI: 10.1016/j.ctim.2017.12.001
- S54. Picelli A, Melotti C, Origano F, Waldner A, Gimigliano R, Smania N. Does robotic gait training improve balance in Parkinson's disease? A randomized controlled trial. *Parkinsonism Relat Disord* 2012; 18(8): 990–993. DOI: 10.1016/j.parkreldis.2012.05.010
- S55. Picelli A, Melotti C, Origano F, Neri R, Waldner A, Smania N. Robot-assisted gait training versus equal intensity treadmill training in patients with mild to moderate Parkinson's disease: A randomized controlled trial. *Park Relat Disord* 2013; 19(6): 605–610. DOI: 10.1016/j.parkreldis.2013.02.010
- S56. Picelli A, Melotti C, Origano F, Neri R, Verze E, Gandolfi M, et al. Robot-assisted gait training is not superior to balance training for improving postural instability in patients with mild to moderate Parkinson's disease: a single-blind randomized controlled trial [with consumer summary]. *Clin Rehabil* 2015; 29(4): 339–347. DOI: 10.1177/0269215514544041
- S57. Romenets SR, Anang J, Fereshtehnejad S-M, Pelletier A, Postuma R, Rios Romenets S, et al. Tango for treatment of motor and non-motor manifestations in Parkinson's disease: A randomized control study. *Complement Ther Med* 2015; 23(2): 175–184. DOI: 10.1016/j.ctim.2015.01.015
- S58. Sage MD, Almeida QJ. Symptom and gait changes after sensory attention focused exercise vs aerobic training in Parkinson's disease. *Mov Disord* 2009; 24(8): 1132–1138. DOI: 10.1002/mds.22469
- S59. Santos SM, da Silva RA, Terra MB, Almeida IA, de Melo LB, Ferraz HB, et al. Balance versus resistance training on postural control in patients with Parkinson's disease. A randomised controlled trial. *Eur J Phys Rehabil Med* 2017; 53(2): 173–183. DOI: 10.23736/S1973-9087.16.04313-6
- S60. Schlenstedt C, Paschen S, Kruse A, Raethjen J, Weisser B, Deuschl G. Resistance versus balance training to improve postural control in Parkinson's disease: A randomized rater blinded controlled study. *PLoS One* [Serial on the Internet] 2015; 10(10). DOI: 10.1371/journal.pone.0140584
- S61. Sedaghati P, Daneshmandi H, Karimi N, Barati A-H. A Selective Corrective Exercise to Decrease Falling and Improve Functional Balance in Idiopathic Parkinson's Disease. *Trauma Mon* 2016; 21(1): e23573. DOI: 10.5812/traumamon.23573
- S62. Shen X, Mak MKY. Balance and Gait Training With Augmented Feedback Improves Balance Confidence in People With Parkinson's Disease: A Randomized Controlled Trial. *Neurorehabil Neural Repair* 2014; 28(6): 524–535. DOI: 10.1177/1545968313517752
- S63. Silva-Batista C, Corcos DM, Roschel H, Kanegusuku H, Gobbi LTB, Piemonte MEP, et al. Resistance training with instability for patients with Parkinson's disease. *Med Sci Sports Exerc* 2016; 48(9): 1678–1687. DOI: 10.1249/MSS.0000000000000945
- S64. Silva-Batista C, Corcos DM, Kanegusuku H, Piemonte MEP, Gobbi LTB, de Lima-Pardini AC, et al. Balance and fear of falling in subjects with Parkinson's disease is improved after exercises with motor complexity. *Gait Posture* 2018; 61: 90–97. DOI: 10.1016/j.gaitpost.2017.12.027
- S65. Smania N, Corato E, Tinazzi M, Stanzani C, Fiaschi A, Girardi P, et al. Effect of balance training on postural instability in patients with idiopathic parkinson's disease. *Neurorehabil Neural Repair* [Serial on the Internet] 2010; 24(9): 826–834. DOI: 10.1177/1545968310376057

- S66. Soke F, Guclu-Gunduz A, Kocer B, Fidan I, Keskinoglu P. Task-oriented circuit training combined with aerobic training improves motor performance and balance in people with Parkinson's Disease. *Acta Neurol Belg* 2021; 121(2): 535–543. DOI: 10.1007/s13760-019-01247-8
- S67. Solla P, Cugusi L, Bertoli M, Cereatti A, Della Croce U, Pani D, et al. Sardinian Folk Dance for Individuals with Parkinson's Disease: A Randomized Controlled Pilot Trial. *J Altern Complement Med* 2019; 25(3): 305–316. DOI: 10.1089/acm.2018.0413
- S68. Song J, Paul SS, Caetano MJD, Smith S, Dibble LE, Love R, et al. Home-based step training using videogame technology in people with Parkinson's disease: a single-blinded randomised controlled trial. *Clin Rehabil* 2018; 32(3): 299–311. DOI: 10.1177/0269215517721593
- S69. Sparrow D, DeAngelis TR, Hendron K, Thomas CA, Saint-Hilaire M, Ellis T. Highly Challenging Balance Program Reduces Fall Rate in Parkinson Disease. *J Neurol Phys Ther* 2016; 40(1): 24–30. DOI: 10.1097/NPT.0000000000000111
- S70. Szeffler-Derela J, Arkuszewski M, Knapik A, Wasiuk-Zowada D, Gorzkowska A, Krzystanek E. Effectiveness of 6-week nordic walking training on functional performance, gait quality, and quality of life in Parkinson's disease. *Med* 2020; 56(7): 1–10. DOI: 10.3390/medicina56070356
- S71. Tollár J, Nagy F, Kovács N, Hortobágyi T. A High-Intensity Multicomponent Agility Intervention Improves Parkinson Patients' Clinical and Motor Symptoms. *Arch Phys Med Rehabil* 2018; 99(12): 2478–2484.e1. DOI: 10.1016/j.apmr.2018.05.007
- S72. Tollár J, Nagy F, Kovács N, Hortobágyi T, Tollár J, Nagy F, et al. Two-Year Agility Maintenance Training Slows the Progression of Parkinsonian Symptoms. *Med Sci Sports Exerc* 2019; 51(2): 237–245. DOI: 10.1249/MSS.0000000000001793
- S73. Tollár J, Nagy F, Hortobágyi T, Tollár J, Nagy F, Hortobágyi T, et al. Vastly Different Exercise Programs Similarly Improve Parkinsonian Symptoms: A Randomized Clinical Trial. *Gerontology [Serial on the Internet]* 2019; 65(2): 120–127. DOI: 10.1159/000493127
- S74. van der Kolk NM, de Vries NM, Kessels RPC, Joosten H, Zwinderman AH, Post B, et al. Effectiveness of home-based and remotely supervised aerobic exercise in Parkinson's disease: a double-blind, randomised controlled trial. *Lancet Neurol* 2019; 18(11): 998–1008. DOI: 10.1016/S1474-4422(19)30285-6
- S75. Van Puymbroeck M, Walter AA, Hawkins BL, Sharp JL, Woschkolup K, Urrea-Mendoza E, et al. Functional Improvements in Parkinson's Disease Following a Randomized Trial of Yoga. *Evid Based Complement Alternat Med* 2018; 2018: 8516351. DOI: 10.1155/2018/8516351
- S76. Vergara-Díaz G, Osypiuk K, Hausdorff JM, Bonato P, Gow BJ, Miranda JG, et al. Tai Chi for Reducing Dual-task Gait Variability, a Potential Mediator of Fall Risk in Parkinson's Disease: A Pilot Randomized Controlled Trial. *Glob Adv Heal Med* 2018; 7: 2164956118775385. DOI: 10.1177/2164956118775385
- S77. Vieira de Moraes Filho A, Chaves SN, Martins WR, Tolentino GP, de Cássia Pereira Pinto Homem R, Landim de Farias G, et al. Progressive resistance training improves bradykinesia, motor symptoms and functional performance in patients with parkinson's disease. *Clin Interv Aging [Serial on the Internet]* 2020; 15: 87-95. DOI: 10.2147/CIA.S231359
- S78. Volpe D, Signorini M, Marchetto A, Lynch T, Morris ME. A comparison of Irish set dancing and exercises for people with Parkinson's disease: a phase II feasibility study. *BMC Geriatr* 2013; 13: 54. DOI: 10.1186/1471-2318-13-54
- S79. Wallén MB, Hagströmer M, Conradsson D, Sorjonen K, Franzén E, Wallén MB, et al. Long-term effects of highly challenging balance training in Parkinson's disease—a randomized controlled trial. *Clin Rehabil* 2018; 32(11): 1520–1529. DOI: 10.1177/0269215518784338
- S80. Wan Z, Liu X, Yang H, Li F, Yu L, Li L, et al. Effects of Health Qigong Exercises on Physical Function on Patients with Parkinson's Disease. *J Multidiscip Healthc* 2021; 14: 941–950. DOI: 10.2147/JMDH.S303945
- S81. Wong-Yu IS, Mak MK. Task- and Context-Specific Balance Training Program Enhances Dynamic Balance and Functional Performance in Parkinsonian Nonfallers: A Randomized Controlled Trial With Six-Month Follow-Up. *Arch Phys Med Rehabil* 2015; 96(12): 2103–2111. DOI: 10.1016/j.apmr.2015.08.409
- S82. Wong-Yu ISK, Mak MKY. Multi-dimensional balance training programme improves balance and gait performance in people with Parkinson's disease: A pragmatic randomized controlled trial with 12-month follow-up. *Parkinsonism Relat Disord* 2015; 21(6): 615–621. DOI: 10.1016/j.parkreldis.2015.03.022
- S83. Xiao C-M, Zhuang Y-C. Effect of health Baduanjin Qigong for mild to moderate Parkinson's disease. *Geriatr Gerontol Int* 2016; 16(8): 911–919. DOI: 10.1111/ggi.12571
- S84. Youm C, Kim Y, Noh B, Lee M, Kim J, Cheon S-M. Impact of trunk resistance and stretching exercise on fall-related factors in patients with Parkinson's disease: a randomized controlled pilot study. *Sensors (Basel)* 2020; 20(15). DOI: 10.3390/s20154106
- S85. Yuan R-Y, Chen S-C, Peng C-W, Lin Y-N, Chang Y-T, Lai C-H. Effects of interactive video-game-based exercise on balance in older adults with mild-to-moderate Parkinson's disease. *J Neuroeng Rehabil* 2020; 17(1): 91. DOI: 10.1186/s12984-020-00725-y
- S86.** Zhang T-Y, Hu Y, Nie Z-Y, Jin R-X, Chen F, Guan Q, et al. Effects of Tai Chi and Multimodal Exercise Training on Movement and Balance Function in Mild to Moderate Idiopathic Parkinson Disease. *Am J Phys Med Rehabil* 2015; 94(10 Suppl 1): 921–929. DOI: 10.1097/PHM.0000000000000351