

Supplementary material to article by G. Francisco et al. “Adult spasticity international registry study: methodology and baseline patient, healthcare provider, and caregiver characteristics”

Table SIII. Spasticity treatment modalities at baseline by lower limb presentation

	Adducted thigh foot (n = 151)	Equinovarus (n = 443)	Flexed hip (n = 59)	Flexed knee (n = 172)	Flexed toe (n = 138)	Stiff extended knee (n = 171)	Striatal/ hyperextended/ hitchhiker toe (n = 57)	Other (n = 36)	Total (n = 616)
Acupuncture, n (%)	150 14 (9.3)	428 42 (9.8)	59 6 (10.2)	169 17 (10.1)	137 13 (9.5)	168 15 (8.9)	55 4 (7.3)	35 6 (17.1)	593 53 (8.9)
Assistive Devices, n (%)	150 115 (76.7)	428 293 (68.5)	59 50 (84.7)	169 137 (81.1)	137 102 (74.5)	168 135 (80.4)	55 40 (72.7)	35 26 (74.3)	593 425 (71.7)
Casting, n (%)	150 9 (6.0)	428 46 (10.7)	59 7 (11.9)	169 11 (6.5)	137 13 (9.5)	168 14 (8.3)	55 4 (7.3)	35 3 (8.6)	593 60 (10.1)
Chemodenervation, n (%)	150 4 (2.7)	428 17 (4.0)	59 1 (1.7)	169 5 (3.0)	137 3 (2.2)	168 12 (7.1)	55 2 (3.6)	35 0 (0.0)	593 25 (4.2)
Intrathecal therapy, n (%)	150 22 (14.7)	428 33 (7.7)	59 11 (18.6)	169 24 (14.2)	137 15 (10.9)	168 19 (11.3)	55 3 (5.5)	35 5 (14.3)	593 61 (10.3)
Orthotics, n (%)	150 78 (52.0)	429 253 (59.0)	59 31 (52.5)	169 91 (53.8)	137 87 (63.5)	168 96 (57.1)	55 31 (56.4)	35 18 (51.4)	594 329 (55.4)
Physio or occupational therapy, n (%)	150 135 (90.0)	429 366 (85.3)	59 53 (89.8)	169 140 (82.8)	137 125 (91.2)	168 148 (88.1)	55 47 (85.5)	35 27 (77.1)	594 491 (82.7)
Surgeries or other procedures, n (%)	150 19 (12.7)	429 53 (12.4)	59 9 (15.3)	169 24 (14.2)	137 16 (11.7)	168 17 (10.1)	55 8 (14.5)	35 4 (11.4)	594 79 (13.3)

NOTE: Allowed responses included “yes” or “not mentioned”.