

Supplemental Table. Summary of patient cohort from subgroups

	2020 ^a	2021 ^b
	N = (%)	N = (%)
Age		
In years, mean (SD)	33 (100)	39 (100)
Strata	57 (21)	56 (20)
18-29	5 (15.2)	6 (15.4)
30-49	6 (18.2)	6 (15.4)
50-64	10 (30.3)	13 (33.3)
65-79	7 (21.2)	10 (25.6)
80+	5 (15.2)	4 (10.3)
Male	23 (69.7)	28 (71.8)
ASA-PS		
1 Normal healthy	11 (33.)	11 (28.2)
2 Mild systemic	9 (27.3)	10 (25.6)
3 Severe systemic	12 (36.4)	18 (46.2)
4 Life threatening	1 (3.0)	0 (0)
Living status		
Home independent	25 (75.8)	39 (100)
Substance dependence	8 (24.2)	14 (35.9)
Trauma mechanism		
Fall	20 (60.6)	24 (61.5)
Motor vehicle	5 (15.2)	1 (2.6)
Pedestrian	-	-
Bicycle	2 (6.1)	4 (10.3)
Electric scooter	0 (0)	3 (7.7)
Sports	0 (0)	1 (2.6)
Violence	2 (6.1)	4 (10.3)
Self-harm	-	-
Other/unknown	4 (12.1)	2 (5.1)
Alcohol at time of injury	8 (24.2)	15 (38.5)
Head Injury Severity Scale		
Minimal	3 (9.1)	4 (10.3)
Mild	9 (27.3)	14 (35.9)
Moderate	12 (36.4)	17 (43.5)
Severe	9 (27.3)	4 (10.3)
High-Energy trauma (yes)	8 (24.2)	9 (23.1)
Isolated TBI	18 (54.5)	24 (61.5)
Treatment TBI		
Conservative	21 (63.6)	34 (87.2)
Any neurosurgical procedure	12 (36.4)	5 (12.8)
Intracranial pressure sensor	9 (27.3)	3 (7.7)
Evacuation of mass lesion	4 (12.1)	4 (10.3)
Discharge destination		
Home	7 (21.2)	13 (33.3)
Local hospital	13 (39.4)	20 (51.3)
Specialized rehabilitation	9 (27.3)	3 (7.7)
Nursing home	3 (9.1)	2 (5.1)
Other	1 (3.0)	1 (2.6)