

Supplementary material to article by J. Kaarre et al. "Predictors for self-reported feeling of depression three months after stroke: a longitudinal cohort study"

**Table SII.** Phi correlation coefficients between independent variables entered into binary logistic regression analyses for the 305 patients with stroke included in the analyses

Variables	Age	Sex	NIHSS score	MoCA score	Living alone before stroke	Diabetes	Prior stroke	Barthel Index
Sex	0.07							
NIHSS score	0.07	0.05						
MoCA score	<b>0.33***</b>	0.01	<b>0.14*</b>					
Living alone before stroke	<b>0.21***</b>	<b>0.17**</b>	<b>0.18**</b>	0.06				
Diabetes	0.03	-0.5	0.11	0.08	0.06			
Prior stroke	0.11	-0.12*	0.06	<b>0.15**</b>	0.03		-0.03	
Barthel Index	<b>0.29***</b>	0.04	<b>0.29**</b>	<b>0.25***</b>	<b>0.23***</b>		0.05	0.10
Stroke type	-0.10	-0.05	0.04	-0.01	-0.10	-0.02	-0.05	-0.03

Statistics: phi correlation coefficient for binary variables and strength of statistical significance where \* $p < 0.05$ , \*\* $p < 0.01$ , and \*\*\* $p < 0.001$ . Bold text indicates significant results. NIHSS: National Institutes of Health Stroke Scale; MoCA: Montreal Cognitive Assessment.